

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|--|-----------|---|
| (51) International Patent Classification ⁶ : A61K 31/355, 31/35 | A1 | (11) International Publication Number: WO 96/19218 (43) International Publication Date: 27 June 1996 (27.06.96) |
| (21) International Application Number: PCT/US95/16848 (22) International Filing Date: 22 December 1995 (22.12.95) (30) Priority Data: 08/362,693 22 December 1994 (22.12.94) US (60) Parent Application or Grant (63) Related by Continuation US 08/362,693 (CIP) Filed on 22 December 1994 (22.12.94) (71) Applicant (for all designated States except US): HENKEL CORPORATION [US/US]; Suite 150, 140 Germantown Pike, Plymouth Meeting, PA 19462 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CLARK, James, P. [US/US]; 40 West Bailey Road, Naperville, IL 60565 (US). DUNKER, Manfred, S. [US/US]; 26 Wildwood Trail, Palos Park, IL 60464 (US). (74) Agent: REA, Teresa, Stanek; Henkel Corporation, Suite 150, 140 Germantown Pike, Plymouth Meeting, PA 19462 (US). | | (81) Designated States: AU, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> |
| (54) Title: TOCOPHEROL AND TOCOTRIENOL COMPOSITIONS (57) Abstract Pharmaceutical compositions comprising natural tocotrienols and natural tocopherols are disclosed. Also disclosed is a method for preventing or treating the harmful effects of high serum cholesterol and/or lipid levels in a mammal, said method comprising orally administering approximately 100 mg to approximately 1000 mg of natural tocotrienols and approximately 100 mg to approximately 1000 mg of tocopherols to a mammal, including humans, in need of such prevention or treatment. | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|---------------------------------------|----|--------------------------|
| AT | Austria | GB | United Kingdom | MR | Mauritania |
| AU | Australia | GE | Georgia | MW | Malawi |
| BB | Barbados | GN | Guinea | NE | Niger |
| BE | Belgium | GR | Greece | NL | Netherlands |
| BF | Burkina Faso | HU | Hungary | NO | Norway |
| BG | Bulgaria | IE | Ireland | NZ | New Zealand |
| BJ | Benin | IT | Italy | PL | Poland |
| BR | Brazil | JP | Japan | PT | Portugal |
| BY | Belarus | KE | Kenya | RO | Romania |
| CA | Canada | KG | Kyrgyzstan | RU | Russian Federation |
| CF | Central African Republic | KP | Democratic People's Republic of Korea | SD | Sudan |
| CG | Congo | KR | Republic of Korea | SE | Sweden |
| CH | Switzerland | KZ | Kazakhstan | SI | Slovenia |
| CI | Côte d'Ivoire | LI | Liechtenstein | SK | Slovakia |
| CM | Cameroon | LK | Sri Lanka | SN | Senegal |
| CN | China | LU | Luxembourg | TD | Chad |
| CS | Czechoslovakia | LV | Latvia | TG | Togo |
| CZ | Czech Republic | MC | Monaco | TJ | Tajikistan |
| DE | Germany | MD | Republic of Moldova | TT | Trinidad and Tobago |
| DK | Denmark | MG | Madagascar | UA | Ukraine |
| ES | Spain | ML | Mali | US | United States of America |
| FI | Finland | MN | Mongolia | UZ | Uzbekistan |
| FR | France | | | VN | Viet Nam |
| GA | Gabon | | | | |

- 1 -

TOCOPHEROL AND TOCOTRIENOL COMPOSITIONS

Field of the Invention

The present invention relates to pharmaceutical compositions comprising a combination of natural tocotrienols and natural tocopherols useful as a dietary supplement which exert a protective action on the cardiovascular system. This invention is also directed to a method for preventing treating hypercholesteremia and/or hyperlipidemia by the administration of natural tocotrienols and natural tocopherols.

Background of the Invention

Vitamin E generally refers to the group of compounds (tocol and tocotrienol derivatives) which exhibit qualitatively the biological activity of alpha-tocopherol. They are fat soluble, closely related chemical compounds found in vegetable oils, such as wheat germ oil, rice oil, soy bean oil and the like. Alpha-tocopherol has the greatest biological activity per unit weight while its isomers beta, gamma, delta, epsilon, zeta and eta-tocopherols have vitamin E activity to a lesser extent. The tocopherols and their esters such as tocopherol acetate, tocopherol palmitate, tocopherol succinate and the like are normally water-insoluble.

Tocotrienols and tocopherols (Vitamin E) are therefore known compounds, U.S. Patent 5,190,618 discloses a process for obtaining tocotrienols from palm oil.

There are many vitamin and multivitamin preparations which include vitamin E as a dietary supplement optionally along with other vitamins and trace elements. For a discussion of these preparations, see, Remington's Pharmaceutical Sciences, Eighteenth Edition (1990), Mack Publishing Company pp. 1008-1009 and Handbook of Nonprescription Drugs, Ninth Edition (1990), published by the American Pharmaceutical Association, pp. 447-527.

Tocotrienols have been implicated in the suppression of cholesterolgenesis and arterial thrombosis by inhibition of platelet thromboxane formation. See, for

-2-

example, the disclosure in U.S. Patent 5,157,132 and the references disclosed therein. Tocopherols (Vitamin E), on the other hand, which comprise alpha, beta, gamma and delta tocopherols are essential for normal reproduction, normal development of muscles, normal resistance of erythrocytes to hemolysis and perhaps most importantly are believed to act as an antioxidant.

High serum levels of lipids and cholesterol in humans are generally considered risk factors in the development of many cardiovascular diseases such as coronary artery disease. Consequently, many therapeutic measures have been proposed to treat hypercholesteremia and hyperlipidemia. One such measure is disclosed in U.S. Patent 5,217,992 which relates to the use of tocotrienol to treat hypercholesteremia, hyperlipidemia and thromboembolic disorders.

Another approach is described in the articles, "Vitamin E Consumption And The Risk Of Coronary Disease In Women" by Stampfer et al., The New England Journal of Medicine, 328: 1444-9 (1993), and "Vitamin E Consumption And The Risk of Coronary Heart Disease In Men" by Rimm et al., The New England Journal of Medicine, 328: 1450-6 (1993), which discloses that oxidation of low-density lipoprotein (LDL) plays a role in atherosclerosis. Thus, the oxidation of LDL increases their incorporation into the arterial intima which is an essential step in atherogenesis.

A variety of dietary and drug regimen have been developed or proposed which would block the oxidative modification of LDL. These regimen usually include the ingestion of Vitamin E alone.

Thus, in the foregoing articles, investigators have studied the effect of taking vitamin E with the risk of coronary disease and observed that the use of vitamin E supplements in middle-aged women was associated with a reduced risk of coronary heart disease. Similarly, an association between a high intake of vitamin E and a lower risk of coronary heart disease was also observed in men.

In another study reported in Lancet, 342: 1379-84 (1993), it was observed that high beta-carotene intake reduced the risk of myocardial infarction. Beta-carotene has also been suggested as useful in reducing undesirable vascular

-3-

events in patients with chronic stable angina. See Gaziano et al., "Beta Carotene Therapy for Chronic Stable Angina," Circulation, 82:III, Abstract No. 0796 (1990).

5 However, a need still exists in the art for a method for protecting a mammal, including humans, from the harmful or adverse effects of high serum cholesterol and/or lipids.

Summary of the Invention

10 The present inventors have surprisingly discovered a method for protecting the mammalian body against high blood levels of cholesterol and/or lipids by the ingestion of natural tocotrienols and natural tocopherols, thereby reducing the risk of the development of cardiovascular disease. The art has never appreciated nor suggested such a method for the use of those natural products.

15 While all natural tocotrienols may be used in the practices of the present invention, tocotrienol obtained from palm oil is preferred because of its low cost and ready availability. The naturally occurring tocotrienols are predominantly the d-isomer.

20 The natural tocopherols useful in the practice of the present invention include all the naturally occurring tocopherols, i.e., alpha, beta, gamma and delta as well as the corresponding esters, such as the succinate.

The present invention is directed to a method for preventing or treating the harmful effects of high serum cholesterol and/or lipid levels in a mammal, including humans.

25 According to the present invention, natural tocotrienols and natural tocopherols are administered preferably daily to exert a protective action on the cardiovascular system of the body against high levels of cholesterol and lipids. Preferably, the active agents are administered in an oral dosage form with each dose typically containing approximately 200 to approximately 400 mg

-4-

tocotrienols and approximately 150 to approximately 500 mg tocopherols per unit dose.

**Detailed Description of the
Preferred Embodiments of the Invention**

5 Disease prevention through management of blood serum lipids and cholesterol is an accepted clinical approach. The present inventors have surprisingly now found a method whereby the body of a mammal, including humans is protected against high levels of cholesterol and/or lipid thereby lowering the risk of cardiovascular disease.

10 Natural tocotrienols and natural tocopherols are derived from vegetable oils. Soy oil is the most widely used source. Sunflower, corn, peanut, rapeseed and cottonseed oils may also be used. Natural tocotrienol and natural tocopherols are very different from that produced by chemical synthesis, i.e., synthetic "vitamin E." While the definition of vitamin E is not consistent, for
15 the purposes of the present invention, vitamin E refers to both tocotrienols and tocopherols.

 Synthetic vitamin E is a mixture of eight different stereoisomers, only one of which is molecularly equivalent to natural vitamin E. The other seven stereoisomers have a lower biological activity. The mammalian body prefers the
20 natural stereoisomer.

 Natural vitamin E is recognized as having 36 percent greater potency than synthetic vitamin E. Recent studies suggest that natural vitamin E is probably twice as effective as synthetic vitamin E.

 Natural vitamin E also remains in the body much longer than synthetic
25 vitamin E. The seven synthetic stereoisomers are secreted into the bile and then into the intestine for removal from the body. The natural vitamin E stereoisomer, on the other hand, is returned to the bloodstream in the form of low density lipoproteins.

-5-

Any natural tocopherol or tocotrienol, its ester or compounds convertible to either tocopherols or their esters are suitable for use in the practice of the present invention.

5 The prior art has failed to appreciate any benefit associated with the administration of tocotrienols and tocopherols to a mammal, including humans, to prevent the harmful effects of high serum cholesterol and lipid levels. Further, the prior art has heretofore never recognized any benefit for such a method using natural tocotrienols and natural tocopherols.

10 According to the present invention, the daily administration of approximately 100 mg to approximately 1000 mg of natural tocopherols and approximately 100 mg to approximately 1000 mg natural tocotrienols, exhibit a protective effect against high levels of cholesterol and lipid thereby preventing or treating the subsequent development of coronary heart disease. The dosage of naturally occurring tocotrienol is preferably in the range of approximately 150
15 mg to approximately 500 mg, more preferably approximately 200 mg to approximately 400 mg. The dosage of naturally occurring tocopherol administered to a mammal, including humans, in need of such prevention or treatment is preferably approximately 150 mg to approximately 500 mg per oral dose. The present inventors' method is cost effective since it avoids expense
20 medication to lower elevated level cholesterol and/or lipid.

The oral compositions of the present invention can be made by conventional compounding procedures known in the pharmaceutical art, that is, by mixing the active substances with edible pharmaceutically acceptable non-toxic inert, solid or liquid carriers and/or excipients suitable for systemic
25 administration and conventionally used in oral dosage forms. Additionally, edible, non-toxic pharmaceutically acceptable stabilizers usually used as stabilizers in oral dosage forms or edible, non-toxic pharmaceutically acceptable salts thereof can be included in the compositions of the present invention. All the above carriers, excipients and stabilizers are intended to include only those

-6-

suitable for oral administration and all are conventional and known to the pharmaceutical compounding art.

To carry out the method of the present invention the natural tocotrienols and the natural tocopherols are formulated with an inert pharmaceutically acceptable carrier, suitable for oral administration. Typical solid dosage forms include for example soft gelatin capsules, hard gelatin capsules, tablets, powders, chewable tablets and the like. Typical liquid dosage forms include emulsions, elixirs and the like. Soft gelatin capsules are preferred in the practice of the present invention. These formulations are prepared in accordance with known techniques in the art. Thus, for example the natural tocotrienols and natural tocopherols are mixed with a fixed oil, such as peanut oil. The resulting mixture is optionally blended and dispensed in gelatin capsules. Thus, the natural tocotrienols and natural tocopherols are preferably combined in the same oral dosage form, however, two separate dosage forms, one of each tocotrienol and tocopherol can be administered in two separate dosage forms.

According to the present invention, the patient ingests the capsule preferably daily to obtain the benefit of the administration of the natural tocotrienols and natural tocopherols.

In order to illustrate the practice of the present invention, the following non-limiting example is provided. It will be appreciated that a vast number of additional compositions fall within the scope of the present invention. The Example is provided by way of illustration only and is not intended to limit the invention in any way.

Example

Approximately 200 mg of natural tocotrienols obtained from palm oil and approximately 200 mg of natural tocopherols are mixed together with approximately 200 ml of peanut oil. The oily mixture is then dispensed in soft gelatin capsule and is ready for administration to a patient.

-7-

What is claimed is:

1. A method for preventing or treating the harmful effects of high serum cholesterol and/or lipid levels in a mammal, said method comprising orally administering approximately 100 mg to approximately 1000 mg of natural
5 tocotrienols and approximately 100 mg to approximately 1000 mg of tocopherols to a mammal in need of such prevention or treatment.
2. The method according to claim 1 wherein the amount of tocotrienol is in the range of approximately 150 mg to approximately 500 mg.
3. The method according to claim 2 wherein the amount of
10 tocotrienol is in the range of approximately 200 mg to approximately 400 mg.
4. The method according to claim 1 wherein the amount of tocopherol is in approximately 150 mg to approximately 500 mg.
5. The method according to claim 1 wherein said tocotrienols are derived from palm oil.
- 15 6. The method according to claim 1 further comprising a pharmaceutically acceptable carrier.
7. An oral pharmaceutical composition for the prevention or treatment of the harmful effects of high serum cholesterol and/or lipid levels in a mammal, said composition comprising approximately 100 mg to approximately
20 1000 mg of natural tocotrienols and approximately 100 mg to approximately 1000 mg natural tocopherols and a pharmaceutically acceptable carrier therefor.

-8-

8. The composition according to claim 7 wherein the amount of tocotrienol is in the range of approximately 100 mg to approximately 500 mg.

9. The composition according to claim 8 wherein the amount of tocotrienol is in the range of approximately 200 mg to approximately 400 mg.

5 10. The composition according to claim 7 wherein the amount of tocopherol is approximately 100 mg to approximately 500 mg.

11. The composition according to claim 7 wherein the oral pharmaceutical composition is a solid dosage form.

12. The composition according to claim 12 wherein the solid dosage
10 form is a capsule.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/16848

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/355, 31/35

US CL :514/456, 458

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/456, 458

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS, MEDLINE- TOCOPHEROLS AND VITAMIN E WITH TOCOTRIENOLS OR PALM OIL FOR LOWERING CHOLESTEROL

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|---|-----------------------|
| A | US, A, 5,393,776 (PEARCE) 28 February 1995, see entire document. | 1-12 |
| Y | Medline Abstracts, issued 1991, Tan, D.T. et al., "Effect Of A Palm-Oil-Vitamin E Concentrate On The Serum And Lipoprotein Lipids In Humans", abstract no. 91189080, Am. J. Clin. Nutr. Volume 53 (4 Suppl), 1027s-1030s. | 1-12 |

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

| | | | |
|-----|---|-----|--|
| *A* | document defining the general state of the art which is not considered to be of particular relevance | *T* | later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| *E* | earlier document published on or after the international filing date | *X* | document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| *L* | document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | *Y* | document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| *O* | document referring to an oral disclosure, use, exhibition or other means | *Z* | document member of the same patent family |
| *P* | document published prior to the international filing date but later than the priority date claimed | | |

Date of the actual completion of the international search

23 APRIL 1996

Date of mailing of the international search report

29 APR 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

M. MOEZIE

Telephone No. (703) 308-1235

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.